

A geometrical approach to the dynamics of spinor condensates I: Hydrodynamics

Ryan Barnett,¹ Daniel Podolsky,^{2,3} and Gil Refael¹

¹*Department of Physics, California Institute of Technology, MC 114-36, Pasadena, California 91125, USA*

²*Department of Physics, University of Toronto, Toronto, Ontario M5S 1A7, Canada and*

³*Department of Physics, Technion, Haifa, 32000, Israel*

(Dated: September 17, 2009)

In this work we derive the equations of motion governing the hydrodynamics of spin- F spinor condensates. We pursue a description based on standard physical variables (total density and superfluid velocity), alongside $2F$ ‘spin-nodes’: unit vectors that describe the spin F state, and also exhibit the point-group symmetry of a spinor condensate’s mean-field ground state. The hydrodynamic equations of motion consist of a mass continuity equation, $2F$ Landau-Lifshitz equations for the spin-nodes, and a modified Euler equation. In particular, we provide a generalization of the Mermin-Ho relation to spin one, and find an analytic solution for the skyrmion texture in the incompressible regime of a spin-half condensate. These results exhibit a beautiful geometrical structure that underlies the dynamics of spinor condensates.

I. INTRODUCTION

A central theme of contemporary atomic physics experiment is the dynamics of Bose-Einstein condensates and other correlated atomic gases. Of particular interest are mixtures of several species such as Fermi-Bose mixtures [1, 2, 3], and bosonic gases with an internal spin degrees of freedom, *i.e.*, spinor condensates. Spin-one and spin-two spinor condensates have been realized as particular hyperfine states of alkali atoms [4, 5, 6]. In addition the trapping and cooling of ^{52}Cr atoms has led to the realization of a spin-three condensate [7]. On the theoretical front, since the initial work of Ohmi and Machida [8] and Ho [9] numerous interesting works have followed that discuss ground states, dynamics, and topological excitations of such systems (see, for instance, a review in [10]).

Recently, it was shown within mean-field theory that the ground states of spinor condensates exhibit a high degree of symmetry. This symmetry is opaque in the standard spinor description of the condensate. On the other hand, the symmetry is transparent in the so-called reciprocal state representation. Here, one uses the fact that the mean field ground state of a spin- F condensate can be described by $2F$ coherent spin states orthogonal to it. Each one of these so-called reciprocal states is fully spin-polarized, pointing along some direction on the unit sphere. Since there are $2F$ such reciprocal states, the ground state is uniquely described (up to an overall phase) in terms of the $2F$ points on the unit sphere [11]. For typical spinor condensate Hamiltonians, these points, which we denote “spin nodes”, form highly symmetric configurations. For instance, an $F = 2$ condensate has a cyclic phase, where the spin nodes are arranged in a tetrahedron, as well as a square phase.

The spin-node description of the ground states of spinor condensates provides an intuitive geometrical description of the state of the condensate. In addition, it provides a parametrization which readily exhibits the hidden point-group symmetries of the state. Despite its appeal, however, this parametrization has not been used to describe the dynamics of spinor condensates. Our goal

in this paper is to provide a complete description of the hydrodynamics of spinor condensates in terms of such spin-nodes.

The description of the dynamics we provide in this paper is hydrodynamic in the sense that it focuses on the low energy dynamics of the system associated with locally conserved quantities, or with the slow elastic deformation of spontaneously broken degrees of freedom. As with any fluid, such a macroscopic hydrodynamic description is highly illuminating and useful. Here we derive such a description using the density, superfluid velocity, and the spin-nodes (the $2F$ vectors on the unit sphere) as our basic degrees of freedom. In addition to the Euler equations, which describe mass, momentum, and energy conservation, we obtain $2F$ Landau-Lifshitz equations for the dynamics of the spin-nodes. Furthermore, our derivation gives a natural generalization of the Mermin-Ho relation which connects the vorticity in a ferromagnetic spinor condensate with the Pontryagin index of the order parameter.

The paper is organized as follows. Sec. II provides general background on spinor condensates, and reviews recent progress on the hydrodynamic description of ferromagnetic condensates [12]. In Sec. III we present the spin node representation of spinor-condensate degrees of freedom, and derive several useful identities within this formalism. In Sec. IV we proceed to obtain hydrodynamic equations for the spin-half condensate, using the spin-node formalism, and find an analytic solution for a skyrmion configuration. In Secs. V and VI, we derive the general hydrodynamic equations of motion for the spin-one condensate, and then for an arbitrary spin- F condensate, which includes a generalization of the Mermin-Ho relation [13].

The treatment of the spin degrees of freedom in this paper is exact, and it accounts for the full geometrical structure of the hydrodynamics of spinor condensates. But the precision of the hydrodynamic description here comes at a price: this formalism becomes increasingly complex as the spin F grows, and, for large F , the analysis of its exact form becomes impractical. Nevertheless,

the equations derived here even for large F become quite useful in their linearized form. In an accompanying paper [14], we show how approximate methods – such as linearizing the equations of motion about mean-field solutions – elucidate the low energy properties of spinor condensates with arbitrary spin in a powerful and elegant way.

II. BACKGROUND

A. Hydrodynamics of spinless BECs

For a single component BEC, it is natural to expect a simple hydrodynamic description in terms of density and flow velocity. We take the time-dependent Gross-Pitaevskii equation (GPE) as our starting point:

$$i\partial_t\psi = -\frac{1}{2}\nabla^2\psi + g\rho\psi \quad (1)$$

where $\psi = \sqrt{\rho}e^{i\theta}$ is the macroscopic wave function, and $\rho = |\psi|^2$ is the density (here and after for notational simplicity we will use scaled units). This equation can be recast into the form of local momentum and mass conservation laws; with the superfluid velocity $\mathbf{v} = \nabla\theta$, one obtains [15]:

$$\partial_t\rho = -\nabla(\rho\mathbf{v}) \ ; \ D_t\mathbf{v} = -\nabla\left(g\rho - \frac{\nabla^2\sqrt{\rho}}{2\sqrt{\rho}}\right) \quad (2)$$

where $D_t = \partial_t + \mathbf{v} \cdot \nabla$ is the material derivative. The first of these is the mass continuity equation, while the second is the Euler equation for a fluid, where a quantum pressure term appears.

B. The hydrodynamics of ferromagnetic BECs and the Mermin-Ho relation

In a series of recent experiments, the quench dynamics of a ferromagnetic spin-one condensate was explored [16, 17, 18, 19]. These experiments motivated Lamacraft to develop a hydrodynamic framework for the ferromagnetic BEC in terms of the superfluid velocity, and the director of its ferromagnetic order $\mathbf{n}$ [12]. This description is particularly illuminating when considering the instabilities of the system. This problem was also theoretically considered in Refs. [20, 21, 22, 23, 24, 25, 26].

The GP Lagrangian density describing such a ferromagnetic spinor condensate is given by

$$\mathcal{L} = i\psi_a^*\partial_t\psi_a - \frac{1}{2}\nabla\psi_a^* \cdot \nabla\psi_a - \frac{1}{2}g\rho^2 - \frac{1}{2}c_2\rho^2m^2 \quad (3)$$

where $a = -F, \dots, F$ is summed over all F_z eigenstates, and

$$\rho = \sum_a \psi_a^*\psi_a \quad \mathbf{m} = \frac{1}{\rho} \sum_{ab} \psi_a^*\mathbf{F}_{ab}\psi_b. \quad (4)$$

Lamacraft's approach assumed an incompressible liquid with a wavefunction restricted to the ferromagnetic phase (assuming large g and c_2)

$$\psi_a = e^{i\theta}\Phi_a(\mathbf{n}) \quad (5)$$

where $\Phi_a(\mathbf{n})$ is the highest eigenstate of $\mathbf{n} \cdot \mathbf{F}$. A substitution of this wavefunction into Eq. (3) yielded the following set of hydrodynamic equations [12]:

$$\begin{aligned} \nabla \cdot \mathbf{v} &= 0 \\ D_t\mathbf{n} &= \frac{1}{2}\mathbf{n} \times \nabla^2\mathbf{n}. \end{aligned} \quad (6)$$

Once the density is eliminated, we notice that the spin dynamics are given by a Landau-Lifshitz equation with the material derivative $D_t = \partial_t + \mathbf{v} \cdot \nabla$. In addition, the vorticity is related to the Pontryagin density by

$$\nabla \times \mathbf{v} = \frac{F}{2}\varepsilon_{\alpha\beta\gamma}n_\alpha(\nabla n_\beta \times \nabla n_\gamma). \quad (7)$$

This identity is widely-known as the Mermin-Ho relation [9, 13]. Among other things, such a relation has important consequences for the topological defects in ferromagnetic condensates [9, 27, 28, 29]. Such hydrodynamic equations were also derived in [30] to describe magnetic properties of quantum hall systems.

Making use of this simple description, Lamacraft showed that the helical configuration of the ferromagnetic condensate [18] is unstable. In general, it is clear that such a geometric description simplifies, at least conceptually, the analysis of spinor-condensate dynamics.

C. General magnetic ground state of spinor condensates, and the reciprocal state representation

A general spin- F spinor-condensate is described by a macroscopic wave-function with $2F+1$ complex components ψ_a ($a = -F, \dots, F$). When quantum fluctuations are unimportant, the condensate dynamics is described by the time-dependent Gross-Pitaevskii equation

$$i\partial_t\psi_a = -\frac{1}{2}\nabla^2\psi_a + \frac{\partial V}{\partial\psi_a^*} \quad (8)$$

where V is the spin-dependent interaction energy. The interaction energy is given by the set of parameters g_S , with $S = 0, 2, \dots, 2F$ describing the two-particle interaction strength in the S total angular momentum channel:

$$V_{\text{int}} = \frac{1}{2} \sum_{S,m} g_S \psi_a^* \psi_b^* \langle ab|Sm \rangle \langle Sm|a'b' \rangle \psi_{a'} \psi_{b'}. \quad (9)$$

In the above, $\langle ab|Sm \rangle$ are Clebsch-Gordan coefficients. Note that this expression can also be written as the expectation value of an operator:

$$V_{\text{int}} = \frac{1}{2} \langle \psi |_1 \langle \psi |_2 \mathcal{V}_{\text{int}} | \psi \rangle_2 | \psi \rangle_1 \quad (10)$$

where

$$\mathcal{V}_{\text{int}} = \sum_{S,m} g_S |Sm\rangle \langle Sm| = \sum_S g_S \mathcal{P}_S. \quad (11)$$

In this expression, $\mathcal{P}_S$ projects into the total spin S scattering channel.

The classical (mean-field) ground states occur for uniform condensates which minimize Eq. (9) for fixed density ρ . This minimization was carried out for $F = 1$ [8, 9], $F = 2$ [31], and $F = 3$ [32, 33] yielding a multitude of magnetic phases, which minimize V_{int} in different regions of $\{g_S\}$ parameter space, only one of which (for every F) is ferromagnetic.

Indeed, quite generally, a spin- F spinor condensate may exhibit several flavors of paramagnetic rather than ferromagnetic behavior in its ground state. For example, a spin-one condensate may exhibit the so-called nematic phase, where $\psi_1 = \psi_{-1} = 0$, and $\psi_0 = 1$. The expectation value of the magnetization for such a state is clearly zero along any direction, $\langle \mathbf{F} \cdot \mathbf{n} \rangle = 0$. But in the absence of a ferromagnetic director, $\mathbf{n}$, can we still describe a spinor condensate's magnetic state geometrically?

Such a geometrical method was put forward in Ref. [11], based on the use of spin-coherent states. A spin-coherent state $|\Phi_{\mathbf{n}}\rangle$ is the eigenvector of the operator $\mathbf{F} \cdot \mathbf{n}$ with the largest eigenvalue. The method of Refs. [11] relies on finding the set of $2F$ spin-coherent states, $\{|\Phi_{\mathbf{n}_i}\rangle\}_{i=1}^{2F}$, which annihilate the ground state of a uniform condensate:

$$\langle \Phi_{\mathbf{n}_i} | \psi_{GS} \rangle = 0. \quad (12)$$

The $2F$ states $|\Phi_{\mathbf{n}_i}\rangle$ provide (up to an overall phase) a unique description of the magnetic spin-state of the condensate at each point in space. Such reciprocal spinors give a natural generalization of the ferromagnetic director to the case of paramagnetic condensates. Instead of the geometrically opaque $2F + 1$ complex numbers ψ_a , it allows a description of the magnetic state in terms of $2F$ unit vectors, $\mathbf{n}_i$, or points on the unit sphere.

In addition to its geometrical transparency, such a description also reveals the highly symmetric nature of the mean-field ground states. All the paramagnetic phases found so far correspond to a spin node configuration which is invariant under point symmetry group operations, and sometimes under a larger symmetry. The nematic phase of the $F = 1$ condensate, for instance, is described by two antipodal spin nodes. $F = 2$ condensates can exhibit a nematic phase as well, but also a phase in which the spin-nodes are the vertices of a square, and a phase with the spin nodes at the vertices of a tetrahedron. Such phases are illustrated in Fig. 1.

The reciprocal-spinor description was so far only utilized to discuss equilibrium properties of spinor condensates. The remarkable geometrical properties and hidden symmetries of the mean-field ground state, however, provide ample motivation for employing the spin nodes to obtain a complete description of the dynamics of spinor




FIG. 1: Possible phases for the spin-one and spin-two condensates. Spin-one condensates have ferromagnetic (1a) and nematic phases (1b) while spin-two condensates have ferromagnetic (2a), uniaxial nematic (2b), square biaxial nematic (2c), and a tetrahedral (2d) phases.

condensates. In the following sections we will develop the tools necessary for such a description, and use them to derive both a hydrodynamic description, as well as small oscillation dynamics near mean-field ground states.

III. SPIN NODE DESCRIPTION OF SPIN- F MAGNETIC STATES

The reciprocal-spinor states so far define the spinor condensate's state only implicitly through Eq. (12). In order to be able to use these variables directly, we must invert the relationship, expressing the spinor condensate Lagrangian directly in terms of these variables. In order to find this direct representation, we first separate the wave function into a piece corresponding to the overall density and phase, and a piece describing the local spin state. We write

$$\psi_a = \psi \chi_a \quad (13)$$

where χ_a is a normalized spin- F spinor

$$\sum_a \chi_a^* \chi_a = \langle \chi | \chi \rangle = 1 \quad (14)$$

and the superfluid density is

$$\rho = |\psi|^2. \quad (15)$$

A. Symmetrized spin-node representation

As discussed above, a spin- F spinor $|\chi\rangle$ can be described by $2F$ reciprocal states. On the other hand, such a state can also be described by a fully symmetrized collection of $2F$ spin-half states. Each spin-half state can be parametrized in terms of two coordinates $\Omega = (\theta, \phi)$ on the unit sphere,

$$|\Omega\rangle = \cos\left(\frac{\theta}{2}\right) e^{i\phi/2} |\uparrow\rangle + \sin\left(\frac{\theta}{2}\right) e^{-i\phi/2} |\downarrow\rangle. \quad (16)$$

In this representation,

$$|\chi\rangle = \frac{1}{\mathcal{N}} \sum_{\{\sigma\}} \left(\prod_{i=1}^{2F} \otimes |\Omega_{\sigma_i}\rangle \right) = \frac{\sqrt{(2F)!}}{\mathcal{N}} |\Omega\rangle \quad (17)$$

where $\mathcal{N}$ is a normalization constant, and the sum over σ runs over the $(2F)!$ permutations of the $2F$ labels for the spin-half parts [34]. In Eq. (17) we also defined $|\Omega\rangle$ as the (unnormalized) sum over permutations of the tensor product.

The properties of the above formulation are most easily understood using the Schwinger Boson construction [35] (for review, see [36]). Schwinger Bosons provide an easy way to construct the Hilbert space of a spin- F spinor state. We define two Schwinger boson creation operators: $\hat{a}^\dagger, \hat{b}^\dagger$. An $\hat{a}^\dagger$ boson adds $1/2$ to both the total spin, and to F_z , whereas a $\hat{b}^\dagger$ boson adds $1/2$ to the total spin, but lowers F_z by half. In this notation

$$\begin{aligned} F_{\text{tot}} &= \frac{1}{2}(\hat{a}^\dagger \hat{a} + \hat{b}^\dagger \hat{b}); \\ F_x &= \frac{1}{2}(\hat{a}^\dagger \hat{b} + \hat{b}^\dagger \hat{a}); \\ F_y &= \frac{1}{2i}(\hat{a}^\dagger \hat{b} - \hat{b}^\dagger \hat{a}); \\ F_z &= \frac{1}{2}(\hat{a}^\dagger \hat{a} - \hat{b}^\dagger \hat{b}). \end{aligned} \quad (18)$$

A spin-half spinor is written as:

$$|\Omega\rangle = u|\uparrow\rangle + v|\downarrow\rangle = (u\hat{a}^\dagger + v\hat{b}^\dagger)|0\rangle$$

with $|0\rangle$ the Schwinger-boson vacuum. Here, u and v can be written in terms of the coordinates on the unit sphere as $u = \cos(\theta/2)e^{i\phi/2}$ and $v = \sin(\theta/2)e^{-i\phi/2}$.

A symmetrized tensor product of $2F$ spins within the SB formalism is simply written as:

$$|\Omega\rangle = |\Omega_1 \dots \Omega_{2F}\rangle = \prod_{i=1}^{2F} (u_i \hat{a}^\dagger + v_i \hat{b}^\dagger) |0\rangle. \quad (19)$$

with u_i and v_i parametrized in terms of θ_i, ϕ_i as shown above. We refer to this collection of the $2F$ spin-half states which construct $|\Omega\rangle$ as “spin nodes”.

If we wish to calculate wavefunction overlaps using the Schwinger Boson formalism, we can use Wick’s theorem to obtain

$$\begin{aligned} \langle \Omega^{(a)} | \Omega^{(b)} \rangle &= \langle 0 | \prod_{i=1}^{2F} (u_i^{(a)*} \hat{a} + v_i^{(a)*} \hat{b}) \prod_{j=1}^{2F} (u_j^{(b)} \hat{a}^\dagger + v_j^{(b)} \hat{b}^\dagger) | 0 \rangle \\ &= \sum_{\{\sigma\}} \prod_{i=1}^{2F} \langle 0 | (u_i^{(a)*} \hat{a} + v_i^{(a)*} \hat{b}) (u_{\sigma_i}^{(b)} \hat{a}^\dagger + v_{\sigma_i}^{(b)} \hat{b}^\dagger) | 0 \rangle \\ &= \sum_{\{\sigma\}} \prod_{i=1}^{2F} \langle \Omega_i^{(a)} | \Omega_{\sigma_i}^{(b)} \rangle. \end{aligned} \quad (20)$$

where σ is a permutation of the $2F$ indices that mark the spin-half parts. This result could have also been obtained directly from Eq. (17). Nevertheless, we find it instructive to demonstrate the simple Schwinger Boson construction to obtain the symmetrized states.

B. Connection between spin nodes and reciprocal spinors

Since the symmetrized spin-node representation can be used to express any spin state directly, it makes a grossly overcomplete basis. Nevertheless, its usefulness arises since it perfectly reflects the spin-nodes formalism of the spinor-condensates ground states [11, 37]. In the following we will introduce the necessary new notation for the spin-node formalism; we summarize the new notation in Appendix A

It is simple to see that a spin-coherent state can be written in terms of Schwinger boson states as

$$\begin{aligned} |(\Omega)^{2F}\rangle &= (u\hat{a}^\dagger + v\hat{b}^\dagger)^{2F} |0\rangle \\ &= |\Omega \dots \Omega\rangle. \end{aligned} \quad (21)$$

Thus a coherent state can be thought of as $2F$ spin-nodes pointing in the same direction. (For a summary of the notation see Appendix A.)

As described in Sec. II C, a reciprocal spinor is a coherent state $|(\Omega_r)^{2F}\rangle = |\Omega_r \dots \Omega_r\rangle$ orthogonal to a given spinor $|\Omega\rangle$. Using the construction in terms of symmetrized spin nodes, we can write an equation to determine the reciprocal spinors for a particular state $|\Omega\rangle = |\Omega_1 \dots \Omega_{2F}\rangle$:

$$\langle (\Omega_r)^{2F} | \Omega \rangle = (2F)! \prod_{i=1}^{2F} \langle \Omega_r | \Omega_i \rangle = 0. \quad (22)$$

This equation has $2F$ solutions, each corresponding to a different term in the product vanishing. That is, the i th solution of Eq. (22) is

$$|(\Omega_r)^{2F}\rangle = |(\Omega_i^t)^{2F}\rangle = |\Omega_i^t \dots \Omega_i^t\rangle, \quad (23)$$

where

$$|\Omega_i^t\rangle = (-v_i^* \hat{a}^\dagger + u_i^* \hat{b}^\dagger) |0\rangle \quad (24)$$

is the time-reversed spinor of $|\Omega_i\rangle$ with $\theta_i^t = \pi - \theta_i$, $\phi_i^t = \phi_i + \pi$. Here we use the fact that a spin-half spinor is orthogonal to its time-reversed counterpart.

The set of reciprocal coherent states is, therefore, just the set of states antipodal to the directions in the spin-node set $\{\Omega_i\}_{i=1}^{2F}$ on the unit sphere. Using this construction, Eq. (19) provides us with a direct expression for a spinor condensate state in terms of its reciprocal vectors.

C. Time derivatives of spinors and spin-nodes

As stated above, our goal in this paper is to extract the dynamics in terms of individual spin nodes $|\Omega_i\rangle$. In order to do so, we must be able to isolate the dynamics of each spin node within $|\Omega\rangle$. Consider the time derivative of $|\Omega\rangle$ which will appear in the GPE. We can express it

as a sum of terms in which the time derivative operates on individual spin nodes:

$$\partial_t |\Omega\rangle = |\partial_t \Omega_1 \Omega_2 \dots\rangle + |\Omega_1 \partial_t \Omega_2 \dots\rangle + \dots \quad (25)$$

The trick that allows us to isolate individual spin nodes consists of taking the inner product of $\partial_t |\Omega\rangle$ with the i th reciprocal-state of $|\Omega\rangle$, which is $|(\Omega_i^t)^{2F}\rangle$. All terms which do not involve a time derivative of $|\Omega_i\rangle$ identically vanish and we are left with the single term

$$\langle (\Omega_i^t)^{2F} | \partial_t |\Omega\rangle = (2F)! \langle \Omega_i^t | \partial_t \Omega_i \rangle \prod_{\substack{j=1 \\ j \neq i}}^{2F} \langle \Omega_i^t | \Omega_j \rangle. \quad (26)$$

We will make extensive use of this method for isolating the dynamics of individual spin nodes in the following sections.

D. Geometrical parametrization of the spin-half components: Moving from $|\Omega_i\rangle$ to $\mathbf{n}_i$

All results above were concerned with breaking a spin- F spinor into its $2F$ spin-half parts, $|\Omega_i\rangle$, and with the correspondence between these spin-half parts and the reciprocal coherent states. We would like, however, to understand the dynamics of spinor condensates not only in terms of the spinors, $|\Omega_i\rangle$, but also in terms of the unit vectors that describe them, $\mathbf{n}_i$, where $|\Omega_i\rangle$ is highest value eigenvector of $\mathbf{F} \cdot \mathbf{n}_i$.

The first step in finding the equations of motion in terms of the spin-directors $\mathbf{n}_i$, is to establish an orthonormal triad $(\mathbf{e}_x, \mathbf{e}_y, \mathbf{n})$ that parameterizes the space on S^2 in the vicinity of $\mathbf{n}_i$. In the following we will only consider a single spin-half part, and therefore we drop the index i .

The first element of the triad is $\mathbf{n}$ itself:

$$\mathbf{n} = 2 \langle \Omega | \mathbf{F} | \Omega \rangle \quad (27)$$

where $\mathbf{F}$ is the spin operator, acting in the spin-half Hilbert space. To complete the triad, we again use the time-reversed spin-half ket, $|\Omega^t\rangle$ where $\langle \Omega^t | \Omega \rangle = 0$. With this, we can construct states pointing in the “ x ” and “ y ” directions with respect to $\mathbf{n}$ as

$$|\Omega_x\rangle = \frac{1}{\sqrt{2}}(|\Omega\rangle + |\Omega^t\rangle) \quad (28)$$

$$|\Omega_y\rangle = \frac{1}{\sqrt{2}}(|\Omega\rangle + i|\Omega^t\rangle). \quad (29)$$

These states allow us to complete the orthonormal triad by defining

$$\mathbf{e}_x = 2 \langle \Omega_x | \mathbf{F} | \Omega_x \rangle, \quad \mathbf{e}_y = 2 \langle \Omega_y | \mathbf{F} | \Omega_y \rangle. \quad (30)$$

From these we can construct

$$\mathbf{e}_\pm = \mathbf{e}_x \pm i\mathbf{e}_y. \quad (31)$$

It is useful to note that $\mathbf{F} \cdot \mathbf{e}_\pm$ act as raising and lowering operators. That is,

$$\mathbf{F} \cdot \mathbf{e}_+ |\Omega\rangle = 0; \quad \mathbf{F} \cdot \mathbf{e}_+ |\Omega^t\rangle = |\Omega\rangle, \quad (32)$$

with similar relations holding for lowering operators.

Note that there is an ambiguity in such coordinate systems since $\mathbf{e}_x$ and $\mathbf{e}_y$ can together be rotated about $\mathbf{n}$ which corresponds to the gauge choice for the spinors. That is, the gauge of a spinor can be changed by $|\Omega\rangle \rightarrow e^{i\lambda} |\Omega\rangle$ without changing its direction $\mathbf{n}$. In general, quantities which are gauge invariant cannot depend on the parameterization of the spin and will only involve the unit vectors $\mathbf{n}_i$. We will adhere to the following gauge convention for parameterizing spin-half spinors:

$$|\Omega\rangle = \cos(\theta/2) e^{i\phi/2} |\uparrow\rangle + \sin(\theta/2) e^{-i\phi/2} |\downarrow\rangle \quad (33)$$

In this gauge-choice it is easy to see that:

$$\mathbf{e}_x = \hat{\theta} \quad \mathbf{e}_y = \hat{\phi} \quad (34)$$

where $\hat{\theta}$ and $\hat{\phi}$ are unit vectors from the spherical coordinate system.

To complete the discussion, we make two observations that will simplify the following analysis. First, we express $\mathbf{F}$ in the basis of our triad as

$$\begin{aligned} \mathbf{F} &= (\mathbf{F} \cdot \mathbf{n})\mathbf{n} + (\mathbf{F} \cdot \mathbf{e}_x)\mathbf{e}_x + (\mathbf{F} \cdot \mathbf{e}_y)\mathbf{e}_y \\ &= (\mathbf{F} \cdot \mathbf{n})\mathbf{n} + \frac{1}{2}(\mathbf{F} \cdot \mathbf{e}_+)\mathbf{e}_- + \frac{1}{2}(\mathbf{F} \cdot \mathbf{e}_-)\mathbf{e}_+ \end{aligned} \quad (35)$$

In addition we note that we can use the spin operator $\mathbf{F}$ as a projection onto $|\Omega\rangle$ and its time-reversed partner $|\Omega^t\rangle$ by

$$|\Omega\rangle \langle \Omega| = \frac{1}{2} + \mathbf{n} \cdot \mathbf{F} \quad (36)$$

and

$$|\Omega^t\rangle \langle \Omega^t| = \frac{1}{2} - \mathbf{n} \cdot \mathbf{F}. \quad (37)$$

E. Derivatives of spin-half spinors in terms of the triad $(\mathbf{e}_x, \mathbf{e}_y, \mathbf{n})$

The relations derived and recalled in the previous sections allow us to also write derivatives of spinors in terms of vector quantities and their derivatives. The terms that we will encounter arise from terms such as the isolated time derivatives in Eq. (26). Let us now find this decomposition in terms of the triad $(\mathbf{e}_x, \mathbf{e}_y, \mathbf{n})$ and its differential forms.

Our goal is thus to find:

$$a_\alpha = i \langle \Omega | \partial_\alpha \Omega \rangle \quad \text{and} \quad \langle \Omega^t | \partial_\alpha \Omega \rangle \quad (38)$$

in terms of $(\mathbf{e}_x, \mathbf{e}_y, \mathbf{n})$ and their derivatives. We define a_α in this form for reasons that will become clear later.

The first object in Eq. (38) can be found by considering the quantity

$$\partial_\alpha(\langle \Omega^t | \mathbf{e}_- \cdot \mathbf{F} | \Omega \rangle) = \partial_\alpha 1 = 0. \quad (39)$$

Allowing the derivative to operate on the bra, the ket, and the vector $\mathbf{e}_-$, we find

$$\langle \partial_\alpha \Omega^t | \Omega^t \rangle + \langle \Omega | \partial_\alpha \Omega \rangle = -\frac{1}{2} \mathbf{e}_+ \cdot \partial_\alpha \mathbf{e}_- \quad (40)$$

On the left-hand side we used the facts that $\mathbf{F} \cdot \mathbf{e}_- | \Omega \rangle = | \Omega^t \rangle$ and $\langle \Omega^t | \mathbf{F} \cdot \mathbf{e}_- = \langle \Omega |$. On the right-hand side we used the fact that $\langle \Omega^t | \mathbf{F} | \Omega \rangle = \frac{1}{2} \mathbf{e}_+$ which can be verified from Eq. (35). It is easy to verify that

$$\begin{aligned} \langle \partial_\alpha \Omega^t | \Omega^t \rangle &= \langle \Omega | \partial_\alpha \Omega \rangle, \\ \mathbf{e}_x \cdot \partial_\alpha \mathbf{e}_y &= -\mathbf{e}_y \cdot \partial_\alpha \mathbf{e}_x \end{aligned}$$

from which we find

$$a_\alpha = \frac{1}{2} \mathbf{e}_y \cdot \partial_\alpha \mathbf{e}_x \quad (41)$$

which is the desired result.

To obtain $\langle \Omega^t | \partial_\alpha \Omega \rangle$ we use a similar trick. Starting with

$$0 = \langle \Omega^t | \Omega^t \rangle \langle \Omega^t | \Omega \rangle = \langle \Omega^t | \frac{1}{2} - \mathbf{n} \cdot \mathbf{F} | \Omega \rangle \quad (42)$$

we find

$$\partial_\alpha(\langle \Omega^t | \frac{1}{2} - \mathbf{n} \cdot \mathbf{F} | \Omega \rangle) = 0. \quad (43)$$

As before, allowing the differentiation to act on the bra, the ket, and $\mathbf{n}$ results in

$$\langle \Omega^t | \partial_\alpha \Omega \rangle = \langle \Omega^t | (\partial_\alpha \mathbf{n}) \cdot \mathbf{F} | \Omega \rangle \quad (44)$$

where the term with $\langle \partial_\alpha \Omega^t |$ vanishes since $(\frac{1}{2} - \mathbf{n} \cdot \mathbf{F}) | \Omega \rangle = 0$. Now, using again the decomposition in Eq. (35), we readily find

$$\langle \Omega^t | \partial_\alpha \Omega \rangle = \frac{1}{2} \mathbf{e}_+ \cdot \partial_\alpha \mathbf{n}. \quad (45)$$

This concludes all the tools we will need for our analysis below. We have found how to directly write a spin- F spinor in terms of its spin-nodes, and extract terms having to do with individual spin nodes out of sums arising, e.g., from differentiation. Furthermore we translated the spin-half representation of $|\Omega_i\rangle$ to a set of $2F$ triad bases $(\mathbf{e}_{ix}, \mathbf{e}_{iy}, \mathbf{n}_i)$ which will allow us to parametrize the spin state geometrically. Appendix ?? summarizes the various notation introduced throughout this section.

IV. HYDRODYNAMICS OF SPIN-HALF CONDENSATES

One of our main goals is to write the exact (mean-field) equations of motion for a spinor condensate in terms of

the spin nodes and the superfluid velocity and density. In this section we achieve this goal for spin-half condensates. The equations of motion can be trivially generalized to general spin- F condensates restricted to the ferromagnetic state, when the spinor $|\chi\rangle$ is restricted to be a coherent spin-state. In this case the equations of motion for the condensate reduce to those we find below.

A. Gross-Pitaevskii Lagrangian

In this section we consider the Gross-Pitaevskii Lagrangian. We begin by writing the Lagrangian in a revealing form, using the representation of the bosonic field which separates the spinor order parameter into a product of a density piece and a spin piece, $\psi_a = \psi \chi_a$. The GP Lagrangian is then:

$$\begin{aligned} \mathcal{L} &= i\psi_a^* \partial_t \psi_a - \frac{1}{2} \nabla \psi_a^* \cdot \nabla \psi_a - V_{\text{int}}. \\ &= i\psi^* \partial_t \psi + \rho a_t - \frac{1}{2} |(-i\nabla - \mathbf{a})\psi|^2 - \frac{1}{2} \rho \Upsilon - V_{\text{int}}. \end{aligned} \quad (46)$$

where V_{int} is the spin-related interaction and $\rho = |\psi|^2$ with ψ a complex field. Eq. (46) defines the spin vector potential:

$$a_t \equiv i \langle \chi | \partial_t \chi \rangle ; \mathbf{a} \equiv i \langle \chi | \nabla \chi \rangle, \quad (47)$$

and the quantity

$$\Upsilon \equiv \langle \partial_\alpha \chi | \partial_\alpha \chi \rangle - \langle \chi | \partial_\alpha \chi \rangle \langle \partial_\alpha \chi | \chi \rangle. \quad (48)$$

An interesting observation is that the quantity Υ for a general spin $F = N/2$ can be identified with the CP^N model from quantum field theory [38]. Notice that there is a $U(1)$ gauge freedom in the density-spin decomposition:

$$\psi \rightarrow e^{i\lambda} \psi, |\chi\rangle \rightarrow e^{-i\lambda} |\chi\rangle.$$

The quantity Υ , however, is gauge independent. We make a gauge choice when we write the normalized $|\chi\rangle$ as in Eq. (17), with $|\Omega\rangle$ written as Eq. (19). This $U(1)$ gauge freedom is also reflected in an ambiguity in the choice of the triad arising from the spin-half parts of $|\chi\rangle \propto |\Omega\rangle$, since for each spin-part $\mathbf{e}_x$ and $\mathbf{e}_y$ can together be rotated about $\mathbf{n}$. The choice of a particular triad is set by the gauge choice. In general, quantities which are gauge invariant cannot depend on the parametrization of the spin, and will only involve the unit vectors $\mathbf{n}_i$. The vector potential can be related to the superfluid velocity by

$$\mathbf{v} = \frac{1}{\rho 2i} (\psi_a^* \nabla \psi_a - \psi_a \nabla \psi_a^*) = \nabla \theta - \mathbf{a} \quad (49)$$

where θ is the argument of ψ . So far we have not used the fact that the spin is $F = 1/2$.

B. Geometric representation of hydrodynamic quantities

Now that we know the quantities of interest in the spinor description of the GP Lagrangian, we can translate them to the hydrodynamic variables of density and magnetization direction.

The most important quantity appearing above is the vector potential as defined in Eq. (47). Following the discussion in Sec. IIID we see that for a spin-half condensate the vector potential is

$$a_\alpha = i \langle \Omega | \partial_\alpha \Omega \rangle = \frac{1}{2} \mathbf{e}_y \cdot \partial_\alpha \mathbf{e}_x. \quad (50)$$

The analogy between $\mathbf{a}$ and the vector potential appearing in the Maxwell equations compels us to consider the antisymmetric field tensor $f_{\alpha\beta} = \partial_\alpha a_\beta - \partial_\beta a_\alpha$. Through a series of manipulations this can be written purely in terms of $\mathbf{n}$

$$\begin{aligned} f_{\alpha\beta} &= \frac{1}{2} \partial_\alpha \mathbf{e}_y \cdot \partial_\beta \mathbf{e}_x - \frac{1}{2} \partial_\beta \mathbf{e}_y \cdot \partial_\alpha \mathbf{e}_x \\ &= \frac{1}{2} (\partial_\alpha \mathbf{e}_y \cdot \mathbf{n}) (\partial_\beta \mathbf{e}_x \cdot \mathbf{n}) - \frac{1}{2} (\partial_\beta \mathbf{e}_y \cdot \mathbf{n}) (\partial_\alpha \mathbf{e}_x \cdot \mathbf{n}) \\ &= \frac{1}{2} (\mathbf{e}_y \cdot \partial_\alpha \mathbf{n}) (\mathbf{e}_x \cdot \partial_\beta \mathbf{n}) - \frac{1}{2} (\mathbf{e}_y \cdot \partial_\beta \mathbf{n}) (\mathbf{e}_x \cdot \partial_\alpha \mathbf{n}) \\ &= \frac{1}{2} (\mathbf{e}_y \times \mathbf{e}_x) \cdot (\partial_\alpha \mathbf{n} \times \partial_\beta \mathbf{n}) \\ &= -\frac{1}{2} \mathbf{n} \cdot (\partial_\alpha \mathbf{n} \times \partial_\beta \mathbf{n}). \end{aligned} \quad (51)$$

Note that in the above we have repeatedly used the fact that $\mathbf{v} \cdot \partial \mathbf{v} = 0$ for any unit vector $\mathbf{v}$. The result is the Pontryagin topological density, which is the object of the celebrated Mermin-Ho relation for spin-half spinors [13, 39].

The only remaining term is the gauge invariant quantity Υ , defined in Eq. (48). For a spin-half state, we find:

$$\begin{aligned} \Upsilon &= \langle \partial_\alpha \Omega | \partial_\alpha \Omega \rangle - \langle \partial_\alpha \Omega | \Omega \rangle \langle \Omega | \partial_\alpha \Omega \rangle \\ &= \langle \partial_\alpha \Omega | \Omega^t \rangle \langle \Omega^t | \partial_\alpha \Omega \rangle = \frac{1}{4} (\mathbf{e}_- \cdot \partial_\alpha \mathbf{n}) (\mathbf{e}_+ \cdot \partial_\alpha \mathbf{n}) \\ &= \frac{1}{4} (\partial_\alpha \mathbf{n}) \cdot (\partial_\alpha \mathbf{n}). \end{aligned} \quad (52)$$

Thus the Υ term signifies the stiffness of the superfluid with respect to magnetic gradients (as opposed to simply $U(1)$ phase gradients). Also, since we identified Υ with the Lagrangian density of a CP^1 model, we now reaffirm its equivalence with the nonlinear sigma model [38].

C. Equations of motion for spin-half condensates

Now that we clarified how the hydrodynamic variables arise in the GP Lagrangian density, we are ready to approach the GP equations of motion. In terms of the original variables, the time-dependent GPE for a spin half

condensate is

$$i \partial_t \psi_a = -\frac{1}{2} \nabla^2 \psi_a + g \rho \psi_a \quad (53)$$

where we note that the interaction energy for this case is

$$V_{\text{int}} = \frac{1}{2} g \rho^2. \quad (54)$$

Following the substitution $\psi_a = \psi \chi_a$, with χ_a the entries of the spin-half spinor $|\chi\rangle$, and contraction with $\langle \chi|$ we find

$$i \partial_t \psi + \psi a_t = \frac{1}{2} (-i \nabla - \mathbf{a})^2 \psi + \frac{1}{2} \Upsilon \psi + g \rho \psi$$

where $a_\alpha = (a_t, \mathbf{a})$ is the vector potential introduced previously. Substituting $\psi = f e^{i\theta}$ and multiplying both sides of the equation by $e^{-i\theta}$ gives

$$\begin{aligned} i \partial_t f - \partial_t \theta f + f a_t &= \frac{1}{2} (-\nabla^2 f - i f \nabla \cdot \mathbf{v} - 2 i \mathbf{v} \cdot \nabla f + f v^2) \\ &\quad + \frac{1}{2} \Upsilon f + g \rho f \end{aligned} \quad (55)$$

where $\mathbf{v} = \nabla \theta - \mathbf{a}$. The imaginary part of this gives

$$\partial_t \rho = -\nabla \cdot (\rho \mathbf{v}) \quad (56)$$

which is a mass conservation equation. On the other hand, taking the real part gives

$$\partial_t \theta + \frac{1}{2} v^2 - a_t = \frac{1}{2} \frac{\nabla^2 f}{f} - \frac{1}{2} \Upsilon - g \rho. \quad (57)$$

We take the gradient of both sides of this equation (using the identity $\nabla(v^2) = 2(\mathbf{v} \cdot \nabla) \mathbf{v} + 2 \mathbf{v} \times (\nabla \times \mathbf{v})$) to get

$$D_t \mathbf{v} = \boldsymbol{\epsilon} + (\mathbf{v} \times \mathbf{b}) - \nabla \left(g \rho + \frac{1}{2} \Upsilon - \frac{\nabla^2 \sqrt{\rho}}{2 \sqrt{\rho}} \right). \quad (58)$$

In this we have defined the “electric” and “magnetic” fields $\boldsymbol{\epsilon}$ and $\mathbf{b}$ in the usual way from the vector potential. That is, $\boldsymbol{\epsilon}_\alpha = f_{\alpha t}$ and $\mathbf{b}_\alpha = (\nabla \times \mathbf{a})_\alpha = \frac{1}{2} \epsilon_{\alpha\beta\gamma} f_{\beta\gamma}$, with α, β and γ indicating space directions, and the f tensor defined below. Also, note that we have used the material derivative $D_t = \partial_t + \mathbf{v} \cdot \nabla$. The “electromagnetic force” appearing in the right-hand-side of the Euler equation is a new feature that is not present in single component condensates. This new type of quantum pressure arises from non-uniform spin textures in spinor condensates.

Now we move on to find the equations describing the spin dynamics. To do this, we contract the GPE with the time reversed spinor $\langle \chi_t |$. This gives

$$\begin{aligned} i \langle \chi_t | D_t \chi \rangle &= -\frac{1}{2} (2 i \mathbf{a} \cdot \langle \chi_t | \nabla \chi \rangle + 2 \frac{\nabla f}{f} \cdot \langle \chi_t | \nabla \chi \rangle \\ &\quad + \langle \chi_t | \nabla^2 \chi \rangle). \end{aligned} \quad (59)$$

Using the spin identities developed in Sec. III the following relations can be derived $\langle \chi_t | \nabla^2 \chi \rangle = \frac{1}{2} \partial_\alpha (\mathbf{e}_+ \cdot \partial_\alpha \mathbf{n})$

and $\langle \partial_\alpha \chi | \chi \rangle \langle \chi_t | \partial_\alpha \chi \rangle = -\frac{1}{4} \partial_\alpha \mathbf{e}_+ \cdot \partial_\alpha \mathbf{n}$. In terms of vectors, the above equation is then

$$\frac{i}{2} \mathbf{e}_+ \cdot D_t \mathbf{n} = -\frac{1}{2} \left(\frac{1}{2} \mathbf{e}_+ \cdot \nabla^2 \mathbf{n} + \frac{\partial_\alpha f}{f} \mathbf{e}_+ \cdot \partial_\alpha \mathbf{n} \right) \quad (60)$$

which can be rewritten as

$$\rho D_t \mathbf{n} = \frac{1}{2} (\mathbf{n} \times \partial_\alpha (\rho \partial_\alpha \mathbf{n})) \quad (61)$$

which is a Landau-Lifshitz equation.

Thus, collecting everything, we can write down a complete set of equations describing the dynamics of a spin half condensate:

$$\begin{aligned} -\partial_t \rho &= \nabla \cdot (\rho \mathbf{v}) \\ \nabla \times \mathbf{v} &= -\mathbf{b} \\ D_t \mathbf{v} &= \mathbf{e} + (\mathbf{v} \times \mathbf{b}) - \nabla \left(g\rho + \frac{1}{8} (\partial_\alpha \mathbf{n})^2 - \frac{\nabla^2 \sqrt{\rho}}{2\sqrt{\rho}} \right) \\ \rho D_t \mathbf{n} &= \frac{1}{2} (\mathbf{n} \times \partial_\alpha (\rho \partial_\alpha \mathbf{n})) \end{aligned}$$

where $\mathbf{e}$ and $\mathbf{b}$ are related to the spin direction through the field tensor

$$f_{\alpha\beta} = \begin{pmatrix} 0 & -\mathbf{e}_x & -\mathbf{e}_y & -\mathbf{e}_z \\ \mathbf{e}_x & 0 & \mathbf{b}_z & -\mathbf{b}_y \\ \mathbf{e}_y & -\mathbf{b}_z & 0 & \mathbf{b}_x \\ \mathbf{e}_z & \mathbf{b}_y & -\mathbf{b}_x & 0 \end{pmatrix} = -\frac{1}{2} \mathbf{n} \cdot (\partial_\alpha \mathbf{n} \times \partial_\beta \mathbf{n}). \quad (62)$$

It is interesting to compare these results with those obtained previously in the incompressible regime [12], Eqns. (6). We find that lifting the incompressibility constraint leads to an Euler equation with effective electric and magnetic fields given by the Mermin-Ho relation. In addition, the superfluid density now enters the Landau-Lifshitz equation.

D. Application: skyrmion texture

As an example of the efficiency of the above hydrodynamic equations of motion, let us consider a specific calculation: skyrmion textures in ferromagnetic condensates. For a standard $U(1)$ vortex, the superfluid velocity close to the vortex core diverges as $1/r$. For a scalar condensate, this causes the superfluid density to be depleted in a small region of order of the coherence length around the core. This can be energetically costly if the condensate is near the incompressible regime. On the other hand, this situation can be circumvented for a spinor condensate. Consider for example, a two component (spin-half) condensate $(\psi_\uparrow, \psi_\downarrow)$, and take the $\downarrow$ component to have a $U(1)$ vortex. Then around the vortex core, the density of $\psi_\downarrow$ can be transferred to the vortex-free $\psi_\uparrow$ keeping the total density across the vortex core finite. This is known as the skyrmion configuration which has been argued to be the relevant topological defect for ferromagnetic condensates [9, 28, 40].




FIG. 2: A skyrmion configuration corresponding to Eq. (66).

Let us now derive the analytic time-independent solution of the equations of motion in the incompressible regime having the skyrmion texture shown in Fig. 2. To this end, we take the incompressible limit [12] of the equations of motion for the spin-half condensate obtained in Sec. IV C. Neglecting z -dependence, these are:

$$\nabla \cdot \mathbf{v} = 0 \quad (63)$$

$$\partial_x v_y - \partial_y v_x = \frac{1}{2} \mathbf{n} \cdot (\partial_x \mathbf{n} \times \partial_y \mathbf{n}) \quad (64)$$

$$D_t \mathbf{n} = \frac{1}{2} (\mathbf{n} \times \nabla^2 \mathbf{n}). \quad (65)$$

With small modifications, these equations can also be shown to describe the dynamics of condensates confined to the ferromagnetic phase of arbitrary spin in the incompressible regime. Our aim is to find stationary solutions of these equations having a skyrmion texture given by [9, 28]

$$\mathbf{n} = (\sin(\beta) \cos(\varphi), \sin(\beta) \sin(\varphi), \cos(\beta)) \quad (66)$$

where φ is the azimuthal angle and β is a function of r which is subject to the boundary conditions $\beta(r=0) = 0$ and $\beta(r=R) = \pi$ where R is a distance far from the skyrmion center. Such a spin configuration is shown in Fig. 2.

Given the form $\mathbf{n}$ in Eq. (66), Eqns. (63,64) can be solved to obtain the velocity profile. One finds

$$\mathbf{v} = \frac{\sin^2(\beta/2)}{r} \hat{\varphi}. \quad (67)$$

Note that the boundary condition $\beta(0) = 0$ suppresses the velocity at the origin which diverges as $1/r$ for the standard $U(1)$ vortex. With the assumption of a static configuration, Eq. (65) reduces to

$$\mathbf{v} \cdot \nabla \mathbf{n} = \frac{1}{2} (\mathbf{n} \times \nabla^2 \mathbf{n}). \quad (68)$$

With the expression for $\mathbf{v}$ in Eq. (67), Eq. (68) leads to the following second order differential equation for β

$$r \left(r \frac{d^2 \beta}{dr^2} + \frac{d\beta}{dr} \right) = \sin(\beta). \quad (69)$$

With the boundary conditions, the solution of this differential equation is

$$\beta(r) = 4 \tan^{-1}(r/R). \quad (70)$$

This expression, along with the velocity in Eq. (67) and the spin direction in Eq. (66) constitute an analytic stationary solution to the equations of motion for the skyrmion configuration.

V. HYDRODYNAMICS OF SPIN-ONE CONDENSATES

A. Geometrical representation of spin one hydrodynamic quantities

Now we move on to considering the more complicated case of the spin-one condensate. The spin-one spinor can be broken down into its two spin-half components and be written as $|\chi\rangle = |\Omega\rangle / \sqrt{\langle\Omega|\Omega\rangle}$ where $|\Omega\rangle = |\Omega_1\Omega_2\rangle$, where we again make use of the large-spin notation defined in Eq. (19). The normalization factor for this case is found to be

$$\begin{aligned} \langle\Omega|\Omega\rangle &= \langle\Omega_1|\Omega_1\rangle \langle\Omega_2|\Omega_2\rangle + \langle\Omega_1|\Omega_2\rangle \langle\Omega_2|\Omega_1\rangle \\ &= \frac{3}{2} + \frac{1}{2} \mathbf{n}_1 \cdot \mathbf{n}_2. \end{aligned} \quad (71)$$

It is also instructive to calculate the spin operator's expectation value. To start we can expand into products of spin-half expectation values

$$\begin{aligned} \langle\Omega|\mathbf{F}|\Omega\rangle &= \langle\Omega_1|\mathbf{F}|\Omega_1\rangle + \langle\Omega_2|\mathbf{F}|\Omega_2\rangle \\ &+ \langle\Omega_1|\Omega_2\rangle \langle\Omega_2|\mathbf{F}|\Omega_1\rangle + \langle\Omega_2|\Omega_1\rangle \langle\Omega_1|\mathbf{F}|\Omega_2\rangle. \end{aligned} \quad (72)$$

Then using the identity in Eq. (36), and the fact that in the factored expression, $\mathbf{F}$ is only acting on spin-half states, we obtain $\langle\Omega|\mathbf{F}|\Omega\rangle = \mathbf{n}_1 + \mathbf{n}_2$. Dividing this by the normalization, we get the spin-one expectation value of the magnetization:

$$\mathbf{m} = \langle\chi|\mathbf{F}|\chi\rangle = 2 \frac{\mathbf{n}_1 + \mathbf{n}_2}{3 + \mathbf{n}_1 \cdot \mathbf{n}_2}. \quad (73)$$

By similar techniques, the vector potential for the spin-one case, with some work, can be written as

$$\begin{aligned} a_\alpha &= i \langle\chi|\partial\chi\rangle = \frac{1}{2} \mathbf{e}_{1y} \cdot \partial_\alpha \mathbf{e}_{1x} + \frac{1}{2} \mathbf{e}_{2y} \cdot \partial_\alpha \mathbf{e}_{2x} \\ &+ \frac{1}{2} \frac{(\mathbf{n}_2 \times \mathbf{n}_1) \cdot \partial_\alpha \mathbf{n}_1 + (\mathbf{n}_1 \times \mathbf{n}_2) \cdot \partial_\alpha \mathbf{n}_2}{3 + \mathbf{n}_1 \cdot \mathbf{n}_2}. \end{aligned} \quad (74)$$

One sees that the first two terms in this expression are the vector potentials from the individual spin-half components while the final term, which is gauge invariant,

describes their coupling. This expression was previously obtained in Refs. [41] and [42], where a geometrical relation for the Berry phase of a spin-one spinor was given. The field tensor corresponding to this vector potential can also be similarly computed. The most simplified form we find is

$$\begin{aligned} f_{\alpha\beta} &= \frac{-2}{(3 + \mathbf{n}_1 \cdot \mathbf{n}_2)^2} \times \\ &(2\mathbf{n}_1 \cdot (\partial_\alpha \mathbf{n}_1 \times \partial_\beta \mathbf{n}_1) + 2\mathbf{n}_2 \cdot (\partial_\alpha \mathbf{n}_2 \times \partial_\beta \mathbf{n}_2) \\ &+ (\mathbf{n}_1 + \mathbf{n}_2) \cdot (\partial_\alpha \mathbf{n}_1 \times \partial_\beta \mathbf{n}_2 + \partial_\alpha \mathbf{n}_2 \times \partial_\beta \mathbf{n}_1)). \end{aligned} \quad (75)$$

This is a generalization of the Mermin-Ho relation to the spin-one case. To our knowledge such an expression has not been previously derived. While its geometrical interpretation is not as immediate as the spin-half case (which is the Pontryagin density), this expression might be of use in computing topological invariant quantities for spin-one fields. This formula has a simplified form when locally restricted to mean-field ground states. For instance for the ferromagnetic state ($\mathbf{n} \equiv \mathbf{n}_1 = \mathbf{n}_2$) the above expression reduces to

$$f_{\alpha\beta} = -\mathbf{n} \cdot (\partial_\alpha \mathbf{n} \times \partial_\beta \mathbf{n}). \quad (76)$$

It is also useful to note that for the nematic state ($\mathbf{n} \equiv \mathbf{n}_1 = -\mathbf{n}_2$) the field tensor identically vanishes, $f_{\alpha\beta} = 0$.

Finally, the gauge invariant quantity Υ can be worked out to be

$$\begin{aligned} \Upsilon &= \frac{2}{(3 + \mathbf{n}_1 \cdot \mathbf{n}_2)^2} (\partial_\alpha \mathbf{n}_1 \cdot \partial_\alpha \mathbf{n}_1 + \partial_\alpha \mathbf{n}_2 \cdot \partial_\alpha \mathbf{n}_2 + \partial_\alpha \mathbf{n}_1 \cdot \partial_\alpha \mathbf{n}_2 \\ &+ \mathbf{n}_1 \cdot \mathbf{n}_2 \partial_\alpha \mathbf{n}_1 \cdot \partial_\alpha \mathbf{n}_2 - \mathbf{n}_1 \cdot \partial_\alpha \mathbf{n}_2 \mathbf{n}_2 \cdot \partial_\alpha \mathbf{n}_1). \end{aligned} \quad (77)$$

This is an explicit representation of the CP^2 model which can be viewed as a generalization of the nonlinear sigma model. Here, too, it is instructive to consider what this expression reduces to when locally restricting to mean-field ground states. For the ferromagnetic state, one finds

$$\Upsilon = \frac{1}{2} \partial_\alpha \mathbf{n} \cdot \partial_\alpha \mathbf{n}. \quad (78)$$

On the other hand, for the nematic state Υ reduces to

$$\Upsilon = \frac{1}{4} \partial_\alpha \mathbf{n} \cdot \partial_\alpha \mathbf{n}. \quad (79)$$

B. Spin-one condensate equations of motion

We now proceed to do a similar analysis for the spin one problem. For this we note that the spin one GP energy functional has the form

$$V_{\text{int}} = \frac{1}{2} g \rho^2 + \frac{1}{2} c_2 \rho^2 \mathbf{m}^2 \quad (80)$$

where $\mathbf{m}$ is the expectation value of the spin-one operator. The first two hydrodynamic equations – the

mass continuity equation and the modified Euler equation – are obtained, as before, by contracting the Gross Pitaevskii equation with $\langle \chi |$. The analysis proceeds along similar lines as the spin half case. However, for this case we need the generalization of the Mermin-Ho relation for spin one given in Eq. (75) to give the field tensor and thus the effective electric and magnetic fields, in addition to the spin one expressions for Υ Eq. (77) and the magnetization $\mathbf{m}$, in Eq. (73). With these quantities, the first two equations of motion are

$$\partial_t \rho = -\nabla \cdot (\rho \mathbf{v}) \quad (81)$$

and

$$D_t \mathbf{v} = \boldsymbol{\epsilon} + (\mathbf{v} \times \mathbf{b}) - \nabla \left(g\rho + c_2 \rho m^2 + \frac{1}{2} \Upsilon - \frac{\nabla^2 \sqrt{\rho}}{2\sqrt{\rho}} \right). \quad (82)$$

Next, let us discuss the spin dynamical equations. These are obtained by contracting the GPE with $\langle \Omega_1^t \Omega_1^t |$ and $\langle \Omega_2^t \Omega_2^t |$. As before, this causes several terms to vanish since these spinors are orthogonal to $|\chi\rangle$. Contracting with $\langle \Omega_1^t \Omega_1^t |$ gives the following equation which gives the time derivative of the first node

$$i\mathbf{e}_{1+} \cdot D_t \mathbf{n}_1 = -\frac{1}{2} \Gamma_\alpha^{12} \mathbf{e}_{1+} \cdot \partial_\alpha \mathbf{n}_1 - \frac{1}{2} \mathbf{e}_{1+} \cdot \nabla^2 \mathbf{n}_1 + c_2 \rho \mathbf{e}_{1+} \cdot \mathbf{m}. \quad (83)$$

A similar equation for the time derivative of $\mathbf{n}_2$ is obtained by contracting with $\langle \Omega_2^t \Omega_2^t |$. In the above, we have collected the following terms into the Γ_α^{ij} parameter

$$\begin{aligned} \Gamma_\alpha^{ij} = & 2 \frac{\partial_\alpha \sqrt{\rho}}{\sqrt{\rho}} - \frac{\partial_\alpha (\mathbf{n}_i \cdot \mathbf{n}_j)}{3 + \mathbf{n}_i \cdot \mathbf{n}_j} \\ & - \frac{\mathbf{n}_i \cdot \partial_\alpha \mathbf{n}_j - i \mathbf{n}_i \cdot (\mathbf{n}_j \times \partial_\alpha \mathbf{n}_j)}{1 - \mathbf{n}_i \cdot \mathbf{n}_j} \\ & + i \frac{(\mathbf{n}_j \times \mathbf{n}_i) \cdot \partial_\alpha \mathbf{n}_i + (\mathbf{n}_i \times \mathbf{n}_j) \cdot \partial_\alpha \mathbf{n}_j}{3 + \mathbf{n}_i \cdot \mathbf{n}_j}. \end{aligned} \quad (84)$$

Finally, separating the real and imaginary parts as $\Gamma_\alpha^{ij} = (\Gamma_\alpha^{ij})' + i(\Gamma_\alpha^{ij})''$, we obtain the Landau-Lifshitz equations

$$(D_t + \frac{1}{2}(\Gamma_\alpha^{12})'' \partial_\alpha) \mathbf{n}_1 = \frac{1}{2} \mathbf{n}_1 \times ((\Gamma_\alpha^{12})' \partial_\alpha \mathbf{n}_1 + \nabla^2 \mathbf{n}_1) - c_2 \rho \mathbf{n}_1 \times \mathbf{m} \quad (86)$$

$$(D_t + \frac{1}{2}(\Gamma_\alpha^{21})'' \partial_\alpha) \mathbf{n}_2 = \frac{1}{2} \mathbf{n}_2 \times ((\Gamma_\alpha^{21})' \partial_\alpha \mathbf{n}_2 + \nabla^2 \mathbf{n}_2) - c_2 \rho \mathbf{n}_2 \times \mathbf{m} \quad (87)$$

This provides a complete set of equations describing the dynamics of the spin-one condensate.

VI. HYDRODYNAMICS FOR GENERAL SPIN- F CONDENSATES.

Now that we have considered the hydrodynamic equations for spin-half and spin-one condensates in detail, in the following we will consider the general case. The first

two equations of motion, the mass continuity equation and the Euler equation are found, as before, to be

$$\partial_t \rho = -\nabla \cdot (\rho \mathbf{v}) \quad (88)$$

and

$$D_t \mathbf{v} = \boldsymbol{\epsilon} + (\mathbf{v} \times \mathbf{b}) - \nabla \left(\frac{2V_{\text{int}}}{\rho} + \frac{1}{2} \Upsilon - \frac{\nabla^2 \sqrt{\rho}}{2\sqrt{\rho}} \right). \quad (89)$$

The effective electric and magnetic fields again follow from the field tensor $f_{\alpha\beta}$ constructed from $a_\alpha = i \langle \chi | \partial_\alpha \chi \rangle$. For a general spin, however, such quantities are cumbersome to express directly in terms of the spin nodes, and we will refrain from doing so.

To obtain the Landau-Lifshitz equations, we contract the GPE with $\langle (\Omega_i^t)^{2F} |$. Doing this gives

$$\begin{aligned} i \langle \Omega_i^t | \partial_t \Omega_i \rangle = & -\partial_\alpha \log \left(\frac{\psi}{\sqrt{\langle \Omega | \Omega \rangle}} \right) \langle \Omega_i^t | \partial_\alpha \Omega_i \rangle \\ & - \frac{1}{2} \langle \Omega_i^t | \nabla^2 \Omega_i \rangle - \langle \Omega_i^t | \partial_\alpha \Omega_i \rangle \sum_{j \neq i} \frac{\langle \Omega_i^t | \partial_\alpha \Omega_j \rangle}{\langle \Omega_i^t | \Omega_j \rangle} \\ & + \frac{\rho}{\lambda_i^* \langle \Omega | \Omega \rangle} \langle (\Omega_i^t)^{2F} | {}_2 \langle \Omega | \mathcal{V}_{\text{int}} | \Omega \rangle {}_2 | \Omega \rangle {}_1. \end{aligned} \quad (90)$$

In this expression, we have used the notation for interaction energy introduced in Eq. (10). In addition we have introduced the quantities λ_i ,

$$\lambda_i = (2F)! \prod_{j \neq i} \langle \Omega_j | \Omega_i^t \rangle. \quad (91)$$

Instead of writing Eq. 90 in terms of vectors as in the previous sections, we will stop at this point. This equation provides a natural starting point in the analysis of the linearized equations of motion which will be developed in the companion paper Ref. [14].

VII. CONCLUSIONS

One of the most striking and surprising features of spinor condensates is the hidden symmetry of their mean field ground states. In this work, we have strived to bring this symmetry to the forefront, and describe hydrodynamics of these condensates using the objects that exhibit the hidden symmetry, namely, the spin-nodes, and reciprocal coherent states.

In this work, we derived the hydrodynamic equations of motion for condensates of general spin, demonstrated their use in the computation of the skyrmion configuration of a ferromagnetic spin-half gas, and generalized the Mermin-Ho relation to spin-one condensates.

A hydrodynamic description tries to capture the low-energy behavior of a continuous medium in terms of conserved and other simple quantities. Therefore, it comes

particularly handy when we consider spinor-condensates close to their mean-field ground state. In a companion paper [14], we will concentrate on small oscillations of the spinor condensate in the vicinity of the mean field-ground state. As we will show, it is there that the hidden point-group symmetry becomes most apparent and accessible. Using the spin-node formalism, and the parametrization of the spin-nodes in terms of a stereographic projection, we will reduce the problem of finding the $2F$ spin-wave eigenmodes to a simple question of decomposing a representation of the appropriate point symmetry group to its irreducible representations. We will also provide a simple recipe that allows the direct extraction of the condensate's spin-wave eigenmodes using the derivatives of the spherical harmonics, coupled with the knowledge of atomic orbital degeneracy lifting under a crystal field.

Acknowledgments

It is a pleasure to acknowledge useful conversations with E. Demler, T.-L. Ho, I. Klich, A. Lamacraft, and especially A. Turner. We would like to acknowledge the hospitality of the KITP, supported by NSF PHY05-51164. We are also grateful for support from the Sherman Fairchild Foundation (RB); the Packard and Sloan Foundations, and the Institute for Quantum Information under NSF grants PHY-0456720 and PHY-0803371 (GR); and CIFAR, NSERC, and CRC (DP).

APPENDIX A: NOTATION

In this Appendix, for convenience, we collect in one place the notation used for the representation of the

spinors. Normalized spinors of arbitrary spin are denoted by $|\chi\rangle$. Non-normalized symmetric combinations of spin-half states are denoted as (with bold fonts)

$$|\mathbf{\Omega}\rangle = |\Omega_1\Omega_2\ldots\Omega_{2F}\rangle. \quad (\text{A1})$$

This is then related to $|\chi\rangle$ by

$$|\chi\rangle = \frac{|\mathbf{\Omega}\rangle}{\sqrt{\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle}}. \quad (\text{A2})$$

Coherent spin states occur when all of the spin-half constituent spins point in the same direction. We denote these by

$$|(\Omega_i)^{2F}\rangle = |\Omega_i\Omega_i\ldots\Omega_i\rangle. \quad (\text{A3})$$

We next define the spin state corresponding to $|\mathbf{\Omega}\rangle$ with its i th component time-reversed. We denote these by

$$|T_i\mathbf{\Omega}\rangle = |\Omega_1\Omega_2\ldots\Omega_i^t\ldots\Omega_{2F}\rangle. \quad (\text{A4})$$

Finally, we define the projection operator $\mathcal{P}$ to be

$$\mathcal{P} = 1 - |\chi\rangle\langle\chi|. \quad (\text{A5})$$

-
- [1] F. Schreck, L. Khaykovich, K. L. Corwin, G. Ferrari, T. Bourdel, J. Cubizolles, and C. Salomon, *Phys. Rev. Lett.* **87**, 080403 (2001).
 - [2] Z. Hadzibabic, C. A. Stan, K. Dieckmann, S. Gupta, M. W. Zwierlein, A. Gorlitz, and W. Ketterle, *Phys. Rev. Lett.* **88**, 160401 (2002).
 - [3] G. Roati, F. Riboli, G. Modugno, and M. Inguscio, *Phys. Rev. Lett.* **89**, 150403 (2002).
 - [4] J. Stenger, S. Inouye, D. M. Stamper-Kurn, H. J. Miesner, A. P. Chikkatur, and W. Ketterle, *Nature* **396**, 345 (1998).
 - [5] H. Schmaljohann, M. Erhard, J. Kronjäger, M. Kottke, S. van Staa, L. Cacciapuoti, J. J. Arlt, K. Bongs, and K. Sengstock, *Phys. Rev. Lett.* **92**, 040402 (2004).
 - [6] M. S. Chang, C. D. Hamley, M. D. Barrett, J. A. Sauer, K. M. Fortier, W. Zhang, L. You, and M. S. Chapman, *Phys. Rev. Lett.* **92**, 140403 (2004).
 - [7] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, *Phys. Rev. Lett.* **94**, 160401 (2005).
 - [8] T. Ohmi and K. Machida, *J. Phys. Soc. Japan* **67**, 1822 (1998).
 - [9] T. L. Ho, *Phys. Rev. Lett.* **81**, 742 (1998).
 - [10] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. S. De, and U. Sen, *Advances in Physics* **56**, 243 (2007).
 - [11] R. Barnett, A. Turner, and E. Demler, *Phys. Rev. Lett.* **97**, 180412 (2006).
 - [12] A. Lamacraft, *Phys. Rev. A* **77**, 063622 (2008).
 - [13] N. Mermin and T.-L. Ho, *Phys. Rev. Lett.* **36**, 594 (1976).
 - [14] R. Barnett, D. Podolsky, and G. Refael, *arxiv:0903.1863*.
 - [15] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, 2002).
 - [16] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, *Nature* **443**, 312 (2006).
 - [17] S. R. Leslie, J. Guzman, M. Vengalattore, J. D. Sau, M. L. Cohen, and D. M. Stamper-Kurn, *arxiv:0806.1553*.
 - [18] M. Vengalattore, S. R. Leslie, J. Guzman, and D. M. Stamper-Kurn, *Phys. Rev. Lett.* **100**, 170403 (2008).
 - [19] M. Vengalattore, J. Guzman, S. Leslie, F. Serwane, and D. M. Stamper-Kurn, *arxiv:0901.3800*.
 - [20] H. Saito, Y. Kawaguchi, and M. Ueda, *Phys. Rev. A* **76**, 043613 (2007).
 - [21] A. Lamacraft, *Phys. Rev. Lett.* **98**, 160404 (2007).

- [22] M. Uhlmann, R. Schützhold, and U. R. Fischer, Phys. Rev. Lett. **99**, 120407 (2007).
- [23] B. Damski and W. H. Zurek, Phys. Rev. Lett. **99**, 130402 (2007).
- [24] S. Mukerjee, C. Xu, and J. E. Moore, Phys. Rev. B **76**, 104519 (2007).
- [25] G. I. Mias, N. R. Cooper, and S. M. Girvin, Phys. Rev. A **77**, 023616 (2008).
- [26] R. W. Cherng, V. Gritsev, D. M. Stamper-Kurn, and E. Demler, Phys. Rev. Lett. **100**, 180404 (2008).
- [27] T. Mizushima, K. Machida, and T. Kita, Phys. Rev. Lett. **89**, 030401 (2002).
- [28] E. J. Mueller, Phys. Rev. A **69**, 033606 (2004).
- [29] T. Mizushima, N. Kobayashi, and K. Machida, Phys. Rev. A **70**, 043613 (2004).
- [30] M. Stone, Phys. Rev. B **53**, 16573 (1996).
- [31] C. V. Ciobanu, S. K. Yip, and T. L. Ho, Phys. Rev. A **61**, 033607 (2000).
- [32] R. Diener and T.-L. Ho, Phys. Rev. Lett. **96**, 190405 (2006).
- [33] L. Santos and T. Pfau, Phys. Rev. Lett. **96**, 190404 (2006).
- [34] E. Majorana, Nuovo Cimento **9**, 43 (1932).
- [35] J. Schwinger, *Quantum Theory of Angular Momentum* (Academic, New York, 1994), edited by L. Biedenharn and H. Van Dam.
- [36] A. Auerbach, *Interacting Electrons and Quantum Magnetism* (Springer, New York, 1994).
- [37] R. Barnett, A. Turner, and E. Demler, Phys. Rev. A **76**, 013605 (2007).
- [38] R. Rajaraman, *Solitons and instantons* (North Holland, 1982).
- [39] R. Kamien, Rev. Mod. Phys. **74**, 953 (2002).
- [40] U. A. Khawaja and H. Stoof, Nature **411**, 918 (2001).
- [41] C. Bouchiat and G. W. Gibbons, J. Physique **49**, 187 (1988).
- [42] J. H. Hannay, J. Phys. A **31**, 53 (1998).